

Detecting supersymmetric Q-balls with neutron stars

Jes Madsen

Institute of Physics and Astronomy,

University of Aarhus,

DK-8000 Århus C, Denmark

e-mail: jesm@dfi.aau.dk

(16 May, 1998)

Supersymmetric Q-balls trapped in neutron stars or white dwarfs may cause the stars to explode. Trapping of Q-balls in neutron stars is shown to be less likely, but trapping in neutron star progenitors more likely than hitherto assumed, making neutron stars very sensitive Q-ball “detectors”. White dwarfs only trap potentially dangerous Q-balls in a narrow parameter range.

12.60.Jv, 97.60.Jd, 98.70.Rz, 98.80.Cq. Accepted for publication in PHYSICS LETTERS B

Supersymmetric Q-balls [1] with non-zero baryon number, Q_B , have been suggested as a possible outcome of early Universe physics, and may be closely related to the formation of net baryon number [2,3]. Should they survive, they would contribute to dark matter in the Universe today [2]. Recently it was suggested that Q-balls trapped by neutron stars could convert the star into supersymmetric phase, thereby reducing the stellar mass until an explosion would set in as the minimal neutron star mass was reached [4,5]. Simulations using general relativistic hydrodynamic codes indicate that such explosions would be observable as mini-supernova events, releasing of order 10^{52} erg/s in electron-antineutrinos and 10^{47} erg/s in keV x -rays [6,7]. The events might also be related to the mysterious cosmic gamma-ray bursts [4,5]. A similar Q-ball trapping scenario was suggested for white dwarf stars [4,5].

The discussion of astrophysical consequences of supersymmetric Q-balls in many ways resemble considerations related to other non-topological solitons carrying baryonic charge, in particular lumps of strange quark matter, also known as quark nuggets [8,9]. The major difference in the context of astrophysical constraints and applications stems from the fact that the mass of most previously considered candidates grows proportionally to the baryon number, whereas the mass of supersymmetric Q-balls grows only as $Q_B^{3/4}$.

Here several arguments developed for strange quark matter [10] and other Q-balls [11] will be generalized to the supersymmetric case. In particular it will be shown that trapping of supersymmetric Q-balls in the stellar precursors of neutron stars is more likely than hitherto assumed, thereby increasing the likelihood of neutron star conversion (in contrast, trapping in the neutron star itself is shown to be significantly suppressed relative to the expectations in [4]). A single Q-ball present in the neutron star interior is sufficient to ultimately transform the whole star. The time scale for this transformation depends on the detailed physics, but is likely to be small compared to the age of the Universe [4]. Thus stars could be very sensitive Q-ball “detectors”.

Other limits previously derived for “ordinary” baryonic Q-balls are not relevant for the supersymmetric counterparts. For instance, electrically charged Q-balls present during the era of Big Bang nucleosynthesis may absorb neutrons and repel protons, thereby reducing the important production of ^4He [12,11]. This effect grows in significance with the total surface area of Q-balls present, but since the supersymmetric Q-balls are very compact relative to, e.g. strange quark matter, a simple calculation shows, that no interesting limits on supersymmetric Q-ball parameters can be derived in this manner.

Q-balls exist in the minimal supersymmetric extension of the standard model as coherent states of squarks, sleptons and Higgs fields. In theories with flat scalar potentials solitons with large baryon number can be stable. For a potential $U(\phi) \sim m^4 = \text{const}$ for large scalar field ϕ , the mass of a supersymmetric Q-ball is approximately given by [5] $M_Q \approx 4\pi\sqrt{2}mQ_B^{3/4}/3 \approx 5924\text{GeV}m_TQ_B^{3/4}$, with $m \approx 0.1\text{--}100$ TeV, and $m_T \equiv m/1\text{TeV}$. A Q-ball is stable relative to nucleons of mass m_n for $M_Q < Q_B m_n$, giving the stability constraint $Q_B > 1.59 \times 10^{15}m_T^4$. The binding energy of a nucleon is $I_n \equiv m_n + M(Q_B) - M(Q_B + 1) \approx m_n[1 - 4729m_TQ_B^{-1/4}] \rightarrow m_n$ for $Q_B \rightarrow \infty$.

The radius of a Q-ball is $R_Q \approx Q_B^{1/4}/\sqrt{2}m \approx 1.40 \times 10^{-17}m_T^{-1}Q_B^{1/4}\text{cm}$, giving a geometrical cross section

$$\sigma_g = \pi R_Q^2 \approx 6.11 \times 10^{-34}m_T^{-2}Q_B^{1/2}\text{cm}^2. \quad (1)$$

Supersymmetric Electrically Charged Solitons (SECS) with a net positive electrical charge in the interior (neutralized by a surrounding cloud of electrons) will interact via electrostatic scattering off matter. The cross section for this will be the Bohr cross section

$$\sigma_B = \pi r_B^2 \approx 10^{-16}\text{cm}^2, \quad (2)$$

unless $R_Q > r_B$, which happens for $Q_B > 2.68 \times 10^{34} m_T^4$. In that case, as well as for Supersymmetric Electrically Neutral Solitons (SENS) the geometrical cross section, Eq. (1) is the relevant one.

Q-balls passing through matter (for example crossing through a star) will be slowed down significantly after removing a total mass comparable to their own. The mass removed by scattering is $M_S = \sigma D$, where $D = \int \rho(x) dx$ is the total column density of the gas. For Q-balls moving one stellar diameter through a star of mass $M_\star$ and radius $R_\star$, the total column density is roughly

$$D \approx 5.0 M_\star / R_\star^2 \approx 2 \times 10^{12} M_\star / M_\odot (R_\odot / R_\star)^2 \text{ g cm}^{-2}, \quad (3)$$

where subscript $\odot$ denotes solar values (the value for D assumes the star to be a polytrope of index 3, which is a reasonable approximation for the applications below, and the Q-ball to be moving on a radial trajectory, which is also a good approximation because of the gravitational attraction from the star).

Thus stopping of a Q-ball requires $M_S > M_Q$. This corresponds to stopping of Q-balls with baryon number below some critical Q_S given by

$$Q_S \approx 5.2 \times 10^{21} m_T^{-4/3} \left(\frac{M_\star}{M_\odot} \right)^{4/3} \left(\frac{R_\star}{R_\odot} \right)^{-8/3} \quad (4)$$

for SECS scattering with the Bohr cross section, and

$$Q_S \approx 1.8 \times 10^{-4} m_T^{-12} \left(\frac{M_\star}{M_\odot} \right)^4 \left(\frac{R_\star}{R_\odot} \right)^{-8} \quad (5)$$

for neutral Q-balls (SENS) and large SECS scattering with a geometrical cross section.

For Q-balls in a monoenergetic distribution the accretion rate on stars is

$$\mathcal{R} = 2\pi G M_\star^2 n_\infty v_\infty^{-1} \frac{R_\star}{M_\star} \left(1 + \frac{v_\infty^2 R_\star}{2 M_\star G} \right), \quad (6)$$

where v_∞ and $n_\infty \equiv \rho_\infty / M_Q$ are the characteristic speed and number densities far from the accreting star. Scaling to typical values for dark matter in the Galaxy we define $v_\infty \equiv v_{250} 250 \text{ km s}^{-1}$ and $\rho_\infty \equiv \rho_{24} 10^{-24} \text{ g cm}^{-3}$ and get

$$\mathcal{R} = 2.20 \times 10^{26} \text{ s}^{-1} Q_B^{-3/4} m_T^{-1} \rho_{24} v_{250}^{-1} \frac{M_\star}{M_\odot} \frac{R_\star}{R_\odot} \left[1 + 0.164 v_{250}^2 \frac{M_\odot}{M_\star} \frac{R_\star}{R_\odot} \right]. \quad (7)$$

Whereas geometrical accretion (second term in Eq. (7)) dominates when Q-balls hit the Earth, leading to a flux of

$$F_{\text{Earth}} = 94 Q_B^{-3/4} m_T^{-1} \rho_{24} v_{250} \text{ cm}^{-2} \text{ s}^{-1} \text{ sterad}^{-1}, \quad (8)$$

accretion on stars is mainly governed by gravitational capture (first term in Eq. (7); the gravitational analog of the Rutherford cross section. In contrast, Ref. [5] apparently assumes geometrical accretion on stars and underestimates the rate of SENS accretion on neutron stars by a factor 10^4). The total number of Q-balls hitting a star in time t is therefore

$$N \approx 6.9 \times 10^{33} \frac{t}{\text{yr}} \rho_{24} v_{250}^{-1} m_T^{-1} Q_B^{-3/4} \frac{M_\star}{M_\odot} \frac{R_\star}{R_\odot}, \quad (9)$$

the total baryon number $Q_{\text{accrete}} = N Q_B$, and the total mass

$$M_{\text{accrete}} \approx 3.7 \times 10^{-20} M_\odot \frac{t}{\text{yr}} \rho_{24} v_{250}^{-1} \frac{M_\star}{M_\odot} \frac{R_\star}{R_\odot}. \quad (10)$$

A star is hit by at least one Q-ball in time t if Q-balls with $Q_B \leq Q_1$ contribute with density ρ_{24} , where

$$Q_1 = 1.3 \times 10^{45} \left(\frac{t}{\text{yr}} \right)^{4/3} \rho_{24}^{4/3} v_{250}^{-4/3} m_T^{-4/3} \left(\frac{M_\star}{M_\odot} \right)^{4/3} \left(\frac{R_\star}{R_\odot} \right)^{4/3}. \quad (11)$$

The equations above are crude approximations in several respects. The true distribution of Q-balls would not be monoenergetic. Velocity peaks in the spectrum would appear as a result of the gravitational relaxation processes during galaxy formation [13,14], and some enhancement (by factors of a few [15]) in the local flux hitting the Earth will

result because of Q-balls gravitationally captured in the Solar System [15,16]. While such effects are very important for detailed studies of the dark matter distribution in our Galaxy, they are also very model dependent and sufficiently small that the simple average flux equations used above suffice for the present purpose.

Ref. [10] considered the trapping of lumps of strange quark matter in neutron stars and especially their stellar progenitors, showing that very strong limits could be placed on their cosmic abundance under the assumption that some pulsars had to be “ordinary” neutron stars rather than converted into quark stars (or conversely, that if strange quark matter is stable, then all “neutron” stars are actually quark stars, since some quark lump pollution of interstellar space is unavoidable, for instance from quark star collisions). In Ref. [11] this argument was generalized to other baryonic Q-balls with mass proportional to baryon number.

Kusenko *et al.* [4,5] consider the even more spectacular possibility that supersymmetric Q-balls being trapped by neutron stars may eat up the neutrons, thereby reducing the neutron star mass below the stability limit of roughly $0.1M_\odot$, leading to an explosion that might be related to gamma-burst events.

Whereas capture in neutron stars is possible in certain ranges of Q-ball parameters, it will be shown below that several effects make trapping in neutron stars less likely than envisaged in [4]. In contrast, trapping in neutron star progenitors will be shown to be much more likely than estimated by Kusenko *et al.*, just like the cases studied in [10,11]. The corresponding trapping of electrically neutral Q-balls (SENS) in white dwarf stars is demonstrated not to take place for most parameters, contrary to the findings in [4]. (Only neutral Q-balls are potentially dangerous to white dwarf survival, as white dwarfs contrary to neutron stars do not possess free neutrons capable of interacting with SECS).

Two aspects were left out of the discussion of Q-ball trapping in neutron stars in [4]. First the fact that Q-balls are extremely relativistic when hitting the star, and second the fact that the crust of a neutron star consists of an ion lattice with a substantial structural resistance, capable of preventing light Q-balls from reaching the central regions of the star.

Concerning the first point, gravitational acceleration causes a typical Q-ball to reach the stellar surface with a velocity $v \approx (2GM_\star/R_\star)^{1/2}$ and a kinetic energy per baryon of $1\text{TeV } m_T Q_B^{-1/4}$ for typical neutron star parameters. This means that Q-balls near the stability limit will experience a highly inelastic collision where the Q-ball is heated and perhaps (partly) disintegrates. Since the binding energy per baryon increases with Q_B , this disintegration is only important for Q_B close to the lower limit for stability.

The second point, Q-balls (or fragments from the disintegration just discussed) being trapped in the crustal lattice of the neutron star, thus being unable to reach the central regions and convert the star into supersymmetric phase, is more important, since the star is only molten in the first few months following formation in a supernova explosion. The column density of the $5 \times 10^{28}\text{g}$ solid crust above the neutron drip region, below which free neutrons are available for absorption in Q-balls, is $D_{\text{crust}} \approx 4 \times 10^{15}\text{g cm}^{-2}$. Only charged Q-balls with $Q_B > 1.3 \times 10^{26} m_T^{-4/3}$ and neutral Q-balls with $Q_B > 2.9 \times 10^9 m_T^{-12}$ penetrate the crust without stopping. Q-balls that are stopped may sink to the interior if gravity $GM_\star M_Q/R_\star^2$ exceeds lattice resistance $\epsilon\sigma_Q$, where the structural energy density for crystalline material in neutron star crusts consisting of ions with mass and charge A , Z , and density ρ is roughly $\epsilon \approx 1 \times 10^{13}\text{ergs cm}^{-3} \rho^{4/3} Z^2 A^{-4/3}$. For an iron lattice at 10^{11}g cm^{-3} , $\epsilon_{28} \equiv \epsilon/10^{28}\text{ergs cm}^{-3} \approx 1$.

Stable neutral Q-balls will ultimately sink to the center because of gravity (unless $Q_B < 4 \times 10^2 m_T^{-12} \epsilon_{28}^4 (R_\star/10\text{km})^8 (M_\odot/M_\star)^4$), but charged Q-balls are trapped in the crust if $Q_B < 6.4 \times 10^{23} m_T^{-4/3} \epsilon_{28}^{4/3} (R_\star/10\text{km})^{8/3} (M_\odot/M_\star)^{4/3}$ and will not pose a threat to the survival of the neutron star.

Contrary to the assumption in [4] the most important trapping mechanism for charged supersymmetric Q-balls over large ranges of Q_B is actually trapping in the supernova progenitor, so that Q-balls are present at the time of neutron star formation. Figure 1 shows the flux of Q-balls reaching the Earth according to Eq. (8) as a function of baryon number for those contributions to galactic halo density where at least one Q-ball is trapped in different phases of stellar evolution (just one Q-ball is needed to start the conversion of the neutron star into supersymmetric phase). The curve marked MS corresponds to Q-ball trapping in a $10M_\odot$ main sequence star of radius $6.3R_\odot$, with a main-sequence lifetime of 5×10^7 years, PMS marks trapping in the post-main-sequence phase just prior to a supernova explosion (estimated as 2 months capture in the $1M_\odot$ stellar core of radius $10^{-2}R_\odot$), and NS denotes trapping in 10^9 years in a $1.4M_\odot$, solid crust neutron star of radius 10 km. The horizontal dashed line would be the corresponding sensitivity to capture during the first few months after the supernova, where the neutron star is molten, and light Q-balls are not trapped in the crust. There will be some lower cut-off in Q_B where Q-balls are destroyed because of the relativistic, inelastic collision with the neutron star, but this limit has not been calculated in detail. Instead, the vertical line in the left part of the figure is the ultimate cut-off, corresponding to the lower limit of Q-ball stability for $m_T = 0.1$. Also marked along the first axis are the lower cut-offs for $m_T = 1, 10$, and 100 . The first axis has been scaled to $Q_B m_T^{4/3}$ so that all flux curves are independent of m_T , except the stability cut-off, and the experimental results (EXP), which happen to be valid for baryon numbers down very close to the stability cut-off for a given choice of m_T . The experimental flux-limits are derived from the discussion in Ref. [17] on the magnetic monopole searches

with the Baksan detector, and should be valid for SENS as well as SECS, whereas the other limits given in Figure 1 are valid for SECS only. These experimental limits may be improved by some 3 orders of magnitude if other monopole search results are properly reinterpreted to be valid for Q-balls, but the astrophysical (stellar) “detectors” are seen to be potentially much more sensitive because of the large surface areas and long integration times.

For comparison Figure 1 also shows the expected flux if Q-balls of a given $Q_B m_T^{4/3}$ were responsible for the dark matter in our Galaxy ($\rho_{24} v_{250} = 1$ in Eq. (8)).

Because of the small geometrical cross section of neutral supersymmetric Q-balls, the methods discussed above are much less sensitive to SENS (c.f. Fig. 2). In fact, SENS pass through main sequence stars without being significantly affected, and only neutron stars are able to stop neutral Q-balls (with $Q_B < 4 \times 10^{35} m_T^{-12}$), except for $m_T < 1$, where it may also happen in white dwarfs and in post-main-sequence stars. Only neutral Q-balls with $Q_B < 10^{12} m_T^{-12}$ are stopped in these systems, so white dwarfs are only endangered by trapped SENS if m_T is small (Q-balls with $Q_B < 1.59 \times 10^{15} m_T^4$ are unstable). This differs from the results of Ref. [4].

Trapping of supersymmetric Q-balls in different stellar types has been studied and related to the suggestion that neutron stars or white dwarfs may explode when transformed into a scalar condensate by the accumulated Q-balls. While accumulation of electrically neutral Q-balls in white dwarfs is important only for a very narrow range of parameters, capture of charged and neutral Q-balls in neutron star progenitors complements the capture in the neutron stars themselves to make neutron stars potentially very efficient Q-ball detectors—many orders of magnitude more flux-sensitive than groundbased experiments. A major uncertainty in the conclusion is the actual lifetime of a neutron star being transformed by the Q-balls. Only preliminary estimates have been given [4], showing that it may be in a range of relevance for cosmic gamma-ray bursts. Another unknown (which would work to increase the effective “capture rate”) is the amount of Q-balls (if any) present in the cloud forming the neutron star progenitors. This depends completely on the prior evolution history of the Galaxy and the interstellar gas. But regardless of the uncertainties, it appears that major windows of supersymmetric Q-ball parameters do provide interesting consequences for neutron star evolution.

ACKNOWLEDGMENTS

This work was supported in part by the Theoretical Astrophysics Center under the Danish National Research Foundation.

-
- [1] S. Coleman, Nucl. Phys. B 262 (1985) 263.
 - [2] A. Kusenko, M. Shaposhnikov, Phys. Lett. B 418 (1998) 46.
 - [3] K. Enqvist, J. McDonald, preprint hep-ph/9711514.
 - [4] A. Kusenko, M. Shaposhnikov, P.G. Tinyakov, I.I. Tkachev, Phys. Lett. B 423 (1998) 104.
 - [5] A. Kusenko, V. Kuzmin, M. Shaposhnikov, P.G. Tinyakov, Phys. Rev. Lett. 80 (1998) 3185.
 - [6] M. Colpi, S.L. Shapiro, S.A. Teukolsky, Astrophys. J. 414 (1993) 717.
 - [7] K. Sumiyoshi, S. Yamada, H. Suzuki, W. Hillebrandt, Astron. Astrophys. 334 (1998) 159.
 - [8] E. Witten, Phys. Rev. D 30 (1984) 272.
 - [9] A. De Rújula, S.L. Glashow, Nature 312 (1984) 734.
 - [10] J. Madsen, Phys. Rev. Lett. 61 (1988) 2909.
 - [11] J. Madsen, Phys. Lett. B 246 (1990) 135.
 - [12] J. Madsen, K. Riisager, Phys. Lett. B 158 (1985) 208.
 - [13] J.R. Ipser, P. Sikivie, Phys. Lett. B 291 (1992) 288.
 - [14] P. Sikivie, I.I. Tkachev, Y. Wang, Phys. Rev. Lett. 75 (1995) 2911.
 - [15] K. Freese, M.S. Turner, Phys. Lett. 123B (1983) 293.
 - [16] S. Dimopoulos, S.L. Glashow, E.M. Prucell, F. Wilczek, Nature 298 (1982) 824.
 - [17] I.A. Belolaptikov *et al.*, preprint astro-ph/9802223.

Figure 1




FIG. 1. Flux of SECS Q-balls hitting the Earth versus Q-ball baryon number (multiplied by $m_T^{4/3}$ to make most curves independent of m_T), for which at least one charged Q-ball is trapped in upper main sequence stars (MS), post-main-sequence stars (PMS), neutron stars (NS), and molten neutron stars (MNS). For comparison limits from detector searches are denoted by EXP. The diagonal curve (DM) is the upper flux limit corresponding to galactic dark matter. For $m_T = 0.1$ SECS capture in neutron stars is governed by Eq. (5); the curves NS.1 and MNS.1. See text for further explanations.

Figure 2




FIG. 2. Flux of SENS Q-balls hitting the Earth versus Q-ball baryon number (multiplied by m_T^{12} to make most curves independent of m_T). Most of the notation as in Fig. 1. The numbers after DM for the dark matter curves indicate the value of m_T . WD-curve shows trapping in a 10^9 year old white dwarf of mass $1M_\odot$, radius $10^{-2}R_\odot$. See text for further explanations.